

## Suitability of Tree Species by Native Plant Community (NPC)

Ecological System	Floodplain Forest				Wet Forest					Forested Rich Peatland						Acid Peatland			
	Northern		Southern		Northern			NW	SO	Northern			NW	SO	Northern				
Floristic Region	FFn57	FFn67	FFs59	FFs68	WFn53	WFn55	WFn64	WFW54	WFs57	FPn62	FPn63	FPn71	FPn72	FPn81	FPn82	FPw63	FPs63	APn80	APn81
Quaking aspen	-				9d	2d	3d	1d									-		
Jack pine																			
Red pine																			
White pine						12d				-			-						3wd
White spruce	-				9d	15d	-	4		-			-				5d		
Northern pin oak																			
Black cherry																			
Red cedar																			
Black oak																			
Shagbark hickory																			
Basswood	3wd		6d			10wd			2wd										
Paper birch	8d				3d	8d	7d	6d	5d	3wd	5wd		3wd				3d		-
Sugar maple			13d						3wd										
Northern red oak																			
Red maple						6wd	6wd												
Bur oak	5wd	6wd	14d			14wd			8wd										
Ironwood																			
Big-toothed aspen																			
White oak																			
Bitternut hickory			12w																
Butternut																			
Silver maple	1w	1w	1w	1w															
American elm	6w	4wd	2w	4wd		11wd	8wd	8w	6w				-						
Green ash	4w	2wd	9w	2wd		5wd			7w										
Box elder	7w	5wd	4w	6wd					10w										
Red elm			11w					10wd	9w										
Hackberry			7w	8wd															
Cottonwood			3w	3w															
Black walnut			10w																
Swamp white oak			5w	7wd															
River birch				5d															
Rock elm			15w																
White cedar					1	4	4			4	1	3d			3d				
Balsam fir	9				4d	9d	9d	5		5d	3d						4d		
Balsam poplar					7d	7d		2											
Black ash	2	3	8		2w	1w	1w	3w	1				-			-			
Yellow birch					8wd	3wd	5wd		4										
Tamarack					-	-	2	7		2	4	2	1	1	1	1	1	2	2
Black spruce					5	13				1	2	1d	2	2d	2d	2	2	1d	1d

## Using the Tables

### Purpose

These tables are intended to help foresters decide which tree species to silviculturally favor or introduce on sites that have been classified using the *Field Guides to the Native Plant Communities of Minnesota*. The tables limit and define the set of trees that a site (NPC) can naturally produce in order to meet broad management objectives. Within that set, the forester is free to select crop trees that should have all of their environmental needs met and find themselves growing among species with which they can compete. The underlying theory is that when trees are suited to their site, they are vigorous. Vigor translates to superior quality, resistance to disease, capacity for natural regeneration, and the ability to withstand fluctuations in climate.

### Suitability Index

Suitability is a mathematical calculation. The data for this calculation come from 4,414 vegetation plots that have been classified as belonging to one of 52 forested NPCs. Two metrics -- commonness and local abundance -- are the elements of suitability.

A plant is "suited" to a NPC when we often find it there. Percent presence was our metric of commonness. Similarly, a plant is "suited" to a NPC when it tends to occur in abundance when present. Mean percent cover-when-present was our metric of local abundance. **The suitability index is the product of percent presence and mean percent cover-when-present.**

**Example:** Of the 4,414 sample plots, 256 were classified as Northern Mesic Hardwood Forest (Mh35). Basswood trees occur in 164 of the 256 plots. Thus, its percent presence as a tree is  $(164/256)*100=64.1\%$ . The mean cover of basswood trees on those 164 plots is 15.0%. Thus, its suitability index is  $64.1*15.0=962$ .

### Climate Shift Calculations

Due to global warming, land managers are bracing for local vegetation shifts to plants whose North American ranges are warmer and drier than their habitat in Minnesota. An analysis of range climate was used to assign and adjust "synecological" scores for our plants with regard to moisture (M) and temperature (H). The scores range from 1 (dry/cool) to 5 (wet/warm). The difference between a plant's individual synecological score and the mean synecological score of its community provides some insight as to whether that plant would benefit or suffer should its local environment become warmer or drier.

**Example:** For each of the 256 Mh35 vegetation plots, the M score of all component plants was summed and averaged to yield a score for each plot. Then the plot scores were summed and averaged to yield an M score for the community, which in this case was 2.3. The adjusted M score for basswood is 2.01, which is drier than 2.3. Thus, we assume that basswood would benefit from a slightly drier conditions. Similarly, the H score for basswood is 4.03, which is substantially warmer than the 2.9 mean for the Mh35 community ... suggesting that basswood would greatly benefit if Mh35 sites get warmer.

### What the Letters Mean -- tree affinity for warmer or drier site conditions

For each tree the a lower-case "w" follows the tree ranking when that tree has a warmer synecological score than the average for the community. A lower-case "d" follows the tree ranking if it has a drier synecological score than the community mean.

Example from Mh35 Community			
Quaking aspen	1d	excellent rating; 1st ranked; favored if habitat gets drier, disfavored if warmer	
White cedar	7	good rating; 7th ranked; disfavored if habitat gets warmer and/or drier	
Sugar maple	13wd	fair rating; 13th ranked; favored if habitat gets warmer and/or drier	
Green ash	14w	poor rating; 14th ranked; favored if habitat gets warmer, disfavored if drier	
American elm	--	very poor rating; occurs in trace amounts; ranking and climate shift not presented	
Jack pine		not known to occur in Mh35, no calculations possible	

### What the Numbers Mean -- trees compared to other trees

The cell numbers indicate a tree's suitability index ranking as compared to all other trees. A tree with the number 1 is considered the species most suited to that NPC.

### What the Colors Mean -- trees compared to all plants

Cell colors indicate whether a tree would be an excellent, good, fair, poor, or very poor choice as a crop tree. A tree's assignment to one of these classes was based upon its suitability index when compared to the index of all other common plants in that community. **Example:** For the Mh35 community, there were 113 plants with >5% presence. The suitability index of each plant was placed in a table and the table sorted to rank plants by their index. The ranking was segregated into 4 groups of 23 plants each and a final class of 21 plants. The group of 23 plants with the highest indices were assigned excellent suitability, the next 23 plants were assigned good suitability, etc. Basswood had the 8<sup>th</sup> highest ranking, placing it in the excellent class along with 22 other plants.



